

# Physical Training and Sports Injury Prevention Guidelines

USACHPPM Injury Prevention

## Apply Ice to Injuries Early to Prevent Re-injury

While cryotherapy has been helpful as a treatment modality affecting swelling, pain, range of motion, etc., the evidence is insufficient to recommend for or against the use of cryotherapy for secondary prevention of injury (or re-injury) or to speed return to activity. It is recommended that randomized, controlled clinical studies be conducted to assess the efficacy of the application of ice after injury as an injury prevention measure.

**Background:** Cryotherapy is the topical application of ice for treatment of acute musculoskeletal injuries. When applied intermittently after injury, ice reduces many of the adverse conditions related to the inflammatory or reactive phase of an acute injury (i.e., pain, prolonged immobilization, and reduced range of motion), all of which may extend recovery time. Studies demonstrate that ice will reduce swelling, inflammation, and pain. Ice placed directly over the injured tissue limits the amount of fluids going into the injured area and slows nerve conduction velocity, both of which serve to decrease pain and improve function. Ice is especially effective in the first 24 to 72 hours after injury onset.

Despite the long history of using cryotherapy to control swelling and pain, there are very few randomized, controlled studies providing evidence to substantiate the effect of cryotherapy alone on measures of secondary prevention of injury (reinjury), return to sport participation, return to full activity, or return to full military duty. Several studies have analyzed cryotherapy combined with other therapeutic modalities (i.e., compression, immobilization, elevation, electrical stimulation, etc). Despite the general acceptance of cryotherapy as an effective intervention, evidence on which to base these conclusions is limited. The review of the literature for the effect of cryotherapy alone on return-to-sport-participation metrics shows that cryotherapy may have a positive effect. However, the relatively poor quality of the studies reviewed is of concern.

**\*\*Information taken from Joint Services Physical Training Injury Prevention Work Group p. 83-166.**

### References:

Airaksinen OV, Kryklund N, Latvala K, Kouri JP, Gronblad M, Kolari P. Efficacy of cold gel for soft tissue injuries: a prospective double-blinded trial. *J Bone Joint Surg Am* 2004; 86A:1101.

Basur RL, Shephard E, Mouzas GL. A cooling method in the treatment of ankle sprains. *Practitioner* 1976;216:708-11.

Bleakley C, McDonough S, MacAuley D. The use of ice in the treatment of acute soft-tissue injury: a systematic review of randomized controlled trials. *Am J Sports Med* 2004;32:251-261.

Cote DJ, Prentice WE, Hooker DN, Shields EW. Comparison of three treatment procedures for minimizing ankle sprain swelling. *Phys Ther* 1998;68:1072-6.

Eston R, Peters D. Effects of cold water immersion on the symptoms of exercise-induced muscle damage. *J Sports Sci* 1999;17:231-8.

Grant AE. Massage with ice (cryokinetics) in the treatment of painful conditions of the musculoskeletal system. *Arch Phys Med Rehab* 1964;45:233-8.

Hayden C. Cryokinetics in an early treatment program. *Phys Ther* 1964;44:990-3.

Hocutt JE, Jaffe R, Rylander CR, Beebe JK. Cryotherapy in ankle sprains. *Am J Sports Med* 1982;10:316-9.

Howatson G, Van Someren KA. Ice massage. Effects on exercise-induced muscle damage. *J Sports Med Phys Fitness* 2003;43:500-5.

Hubbard TJ, Aronson SL, Denegar CR. Does cryotherapy hasten return to participation? a systematic review. *J Athl Training* 2004;39:88-94.

Hubbard TJ, Denegar CR. Does cryotherapy improve outcomes with soft tissue injury? *J Athl Training* 2004;39:278-9.

Isabell WK, Durrant E, Myrer W, Anderson S. The effects of ice massage, ice massage with exercise, and exercise on the prevention and treatment of delayed onset muscle soreness. *J Athl Training* 1992;27:208-17.

Laba E, Roestenburg M. Clinical evaluation of ice therapy for acute ankle sprain injuries. *NZJ Physiother* 1989;17:7-9.

MacAuley D. Do textbooks agree on their advice on ice? *Clin J Sports Med* 2001;11:67-72.

MacAuley DC. Ice Therapy: How good is the evidence? *Int J Sports Med* 2001;22:379-384.

McMaster WC. A literary review on ice therapy in injuries. *Am J Sports Med* 1977;5:124-6.

Ogilvie-Harris DJ, Gilbert M. Treatment modalities for soft tissue injuries of the ankle: a critical review. *Clin J Sport Med* 1995;5:175-86.

Sloan JP, Hain R, Pownall R. Clinical benefits of early cold therapy in accident and emergency following ankle sprain. *Arch Emerg Med* 1989;6:1-6.

Swenson C, Sward L, Karlsson J. Cryotherapy in sports medicine (review article). *Scan J Med Sci Sports* 1996;6:193-200.

Thompson C, Kelsberg G, St. Anna L. Heat or ice for acute ankle sprains? *J Fam Pract* 2003;52:642-3.

Wilkerson G B, Horn-Kingery HM. Treatment of the inversion ankle sprain: comparison of different modes of compression and cryotherapy. *J Orthop Sports Phys Ther* 1993;17:240-6.

Yackzan L, Adams C, Francis KT. The effects of ice massage on delayed muscle soreness. *Am J Sports Med* 1984;12:159-65.

Yanagisawa O, Miyanaga Y, Shiraki H, Shimojo H, Mukai N, Niitsu M, Itai Y. The effects of various therapeutic measures on shoulder range of motion and cross-sectional areas of rotator cuff muscles after baseball pitching. *J Sports Med Phys Fitness* 2003;43:356-66.

Yanagisawa O, Miyanaga Y, Shiraki H, Shimojo H, Mukai N, Niitsu M, Itai Y. The effects of various therapeutic measures on shoulder strength and muscle soreness after baseball pitching. *J Sports Med Phys Fitness* 2003;43:189-201.\*\*Information taken from Joint Services Physical Training Injury Prevention Work Group.